



Name: \_\_\_\_\_

Date of Birth: \_\_\_\_\_

## Repair of Tendon or Nerve

Operation: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

Site: \_\_\_\_\_

Side: \_\_\_\_\_

*A repair of tendons and nerves procedure is the surgical repair of the tendons and / or nerves*

There are risks and complications with any surgical procedure. They include but are not limited to the following.

### General Risks:

- Infection can occur, requiring antibiotics and further treatment.
- Bleeding could occur and may require a return to the operating room. Bleeding is more common if you have been taking blood thinning drugs such as Warfarin, Aspirin, Clopidogrel (Plavix or Iscover) or Dipyridamole (Persantin or Asasantin).
- Small areas of the lung can collapse, increasing the risk of chest infection. This may need antibiotics and physiotherapy.
- Increased risk in obese people of wound infection, chest infection, heart and lung complications, and thrombosis.
- Heart attack or stroke could occur due to the strain on the heart.
- Blood clot in the leg (DVT) causing pain and swelling. In rare cases part of the clot may break off and go to the lungs.
- Death as a result of this procedure is possible.

### Specific risks:

- Numbness associated with the use of tourniquet with nerve and muscle damage at the site where the tourniquet was placed. This may be temporary or permanent.
- Skin death under the tourniquet, which may require further dressings and / or surgery and skin grafting.

- Loss of function of the repaired tendons due to rupture or growth of fibrous bands of tissue. This may require further surgery.
- Deformity and stiffness of the affected joint. This may be permanent and may require further surgery.
- Numbness in the area supplied by the nerve. This may be permanent but can sometimes be treated with further surgery.
- Weakness in the area supplied by the damaged nerve and/ or tendon(s). This may be permanent but can sometimes be treated by further surgery.
- Stiffening, swelling and puffiness of all fingers and joints of the limb with skin changes. This may require further surgery to release the swelling.
- Abnormal pain response to surgery with worsening of pain and disability.
- The surgical cut may cause changes to the sensation and colour of the limb.
- In some people, healing of the wound may be abnormal and the wound can be thickened and red and the scar may be painful.

### Anaesthetic Risks:

There are risks associated with all anaesthetic types (including general, spinal or regional). These risks are relative to each patient's past medical and family history. Your anaesthetist will discuss the specifics of your anaesthetic with you prior to surgery.



Name: \_\_\_\_\_

Date of Birth: \_\_\_\_\_

*I agree that I have discussed the risks and benefits of the aforementioned procedure and have had a chance to discuss the treatment options available to me with Dr John Roe.*

*I request to have the procedure.*

**Patient:**

Name: \_\_\_\_\_

Signature: \_\_\_\_\_

Date: \_\_\_\_\_

**Patient Carer or Advocate:**

Name: \_\_\_\_\_

Relationship: \_\_\_\_\_

Signature: \_\_\_\_\_

Date: \_\_\_\_\_

**Surgeon:** Dr John Roe

Signature: \_\_\_\_\_

Date: \_\_\_\_\_